## NEUROLOGICAL AUTOIMMUNE REACTIVITY SCREEN<sup>™</sup>

CANNER?



Antigens Tested (IgG + IgA Combined, IgM tested separately for each antigen)

and solid by a providence of the second

ARR

141

- Myelin Basic Protein
- Asialoganglioside
- Alpha + Beta Tubulin
- Cerebellar
- Synapsin

2 mL Serum

treatment protocols.

CLINICAL USE:	RECOMMENDED FOR PATIENTS WHO:
<ul> <li>Detection of immune reactivity in pathogenesis of neuro-inflammatory, neuro-autoimmune and neuro- degenerative disorders.</li> </ul>	Have idiopathic neurological conditions such as     Cerebellar ataxia or Peripheral Neuropathy.
<ul> <li>Early detection of autoimmune reactivity of both peripheral and central nervous system.</li> </ul>	<ul> <li>Have Gluten-Reactivity, Dairy Sensitivity and/or enhanced permeability of the Gut and/or Blood- Brain Barrier.</li> </ul>
Assist in monitoring the effectiveness of related	



www.Joincyrex.com

©2017 Cyrex Laboratories, LLC. All Rights Reserved.

## NEUROLOGICAL AUTOIMMUNE REACTIVITY SCREEN<sup>™</sup>

