

CLINICAL SPECIFICATIONS

RICE, WHITE + BROWN, COOKED

Antigen Made From:

Rice, White and Brown, boiled in water until fully cooked

Associated With:

Rice immune reactivity

Known Cross-Reactions: Soy;¹ Wheat, Corn;² Gliadin;³ Millet;^{2,4} Islet Cell¹⁷

Clinical Significance:

One hundred grams of cooked white and brown rice contain 2.38% protein.⁵ Studies on food immune reactivities predominantly use raw food antigens. However, some researchers have noted that heating, or combining, food proteins can change their antigenicity.⁶⁻⁸

This array tests for IgG and IgA food immune reactivity.^{9,10} Equivocal or out-of-range results indicate antibody reactivity to the tested food antigen. We tested 288 blood donor sera against cooked white and brown rice antigens at optimal dilution, 19.8% of these donors were IgG and IgA reactive.

Due to cross-reactivity, possible connections between food antigens and human autoimmunity has been previously suggested because proteins in nature can have a similarity in sequence and structure to certain human tissues.¹¹⁻¹⁴

Data suggests that eliminating foods identified using IgG antibody food testing can play a role in improvement of symptoms.¹⁵ Because certain food components can lead to gut flora changes and gut permeability, eliminating specified food antigens should result in the reduction of antigenic stimuli and the improvement of symptoms.^{15,16}

The results of this food array may be used to develop and implement an immune targeted dietary plan, which includes the avoidance of triggering and known cross-reactive foods. Furthermore, when followed over time, avoidance/prevention treatment plans tailored and supervised by the ordering healthcare professional, may help: (a) repair the gut barrier; and (b) re-establish oral tolerance to the offending food.^{15,16}

References:

1. Lehrer et al. Corn Allergens: IgE antibody reactivity and cross-reactivity with rice, soy, and peanut. *Int Arch Allergy Immunol*, 1999; 118:298-299.
2. Urisu et al. 16-kilodalton rice protein is one of the major allergens in rice grain extract and responsible for cross-allergenicity between cereal grains in the poaceae family. *Int Arch Allergy Immunol*, 1991; 96(3):244-252.
3. Vojdani and Tarash. Cross-reaction between gliadin and different food and tissue antigens, *Food Nutri Sci*, 2013; 4:20-32.
4. Yamada et al. The involvement of rice protein 16KD in cross-allergenicity between antigens in rice, wheat, corn, Japanese millet, Italian millet. [Japanese] *Alerugi*, 1991; 40(12):1485-1492.
5. U.S. Department of Agriculture. <http://ndb.nal.usda.gov/ndb/foods>
6. Sanchez and Fremont. Consequences of heat treatment and processing of food on the structure and allergenicity of component proteins. *Rev Fr Allergol Immunol Clin*, 2003; 43:13-20.
7. Sathe et al. Effects of food processing on the stability of food allergens. *Biotechnol Adv*, 2005; 23:423-429.
8. Vojdani. Detection of IgE, IgG, IgA and IgM antibodies against raw and processed food antigens. *Nutr Metab (Lond)*, 2009; 6: 22. DOI: 10.1186/1743-7075-6-22.
9. Barnes. IgG and IgA antibodies to dietary antigens in food allergy and intolerance. *Clin Exp Allergy*, 1995; 25(Suppl 1):7-9.
10. Mullin et al. Testing for food reactions: the good, the bad, and the ugly. *Nutr Clin Pract*, 2010; 25(2):192-198.
11. Vaishnav et al. Aquaporin 4 molecular mimicry and implications for neuromyelitis optica. *J Neuroimmunol*, 2013; 260: 92-98.
12. Agris et al. Plant DNA topoisomerase 1 is recognized and inhibited by human SCI-70 sera autoantibodies. *Exp Cell Res*, 1990;189:276-279.
13. Lunardi et al. Glycine-rich cell wall proteins act as specific antigen targets in autoimmune and food allergic disorders. *Int Immunol*, 2000; 12(5):647-657.
14. Bullard-Dillard et al. Anti-Sm autoantibodies of systemic lupus erythematosus cross react with dietary plant proteins. *Immunol Invest*, 1992; 21(3):193-202.
15. Cordain et al. Modulation of immune function by dietary lectins in rheumatoid arthritis. *Br J Nutr*, 2000; 83:207-217.
16. Atkinson et al. Food elimination based on IgG antibodies in irritable bowel syndrome: a randomised controlled trial. *Gut*, 2004; 53(10):1459-1464.
17. Kharrazian, et al. Detection of islet cell immune reactivity with low glycemic index foods: is this a concern for type 1 diabetes? *J Diabetes Res*, 2017; 2017:4124967.